arXiv:1107.2558v1 [hep-th] 13 Jul 2011

An Exact Solution to the Quantized Electromagnetic Field in the de Sitter Space

G. Alencar,[1, *] I. Guedes,[2, †] R. R. Landim,[2, ‡] and R.N. Costa Filho[2, §]

[1] *Universidade Estadual do Ceará, Faculdade de Educação,*
Ciências e Letras do Sertão Central-R. Epitácio Pessoa, 2554, 63.900-000 Quixadá, Ceará, Brazil.
[2] *Departamento de Física, Universidade Federal do Ceará,*
Caixa Postal 6030, Campus do Pici, 60455-760 Fortaleza, Ceará, Brazil

In this work we investigate the quantum theory of light propagating in a de Sitter space-time. To do so, we use the method of dynamic invariants to obtain the solution of the time-dependent Schrödinger equation. The quantum behavior of the electromagnetic field in this background is analyzed. The quantum dispersion, the uncertainty product and the correlation between coordinate and momentum is evaluated.

INTRODUCTION

The study of fields in a curved background has been considered along the last decades [1]. The quantization of gravity can be carried out by using the String Theory [2–4]. However, in cosmological scales, gravity can be considered as a classical theory and the fields as propagating waves in the background. Time-dependent backgrounds are used to describe many physical systems yielding interesting results. For instance, in the study of black hole evaporation [5] and the Unruh and Casimir effects [6, 7]. They are also very useful to describe the dynamical evolution of the universe, where the production of particles in cosmological spacetimes has been investigated [8–10].

The quantum effects of a massive scalar field in the de Sitter spacetime from a Schrödinger-picture point of view has been invstigated in Ref. [11]. The authors used an exact linear invariant and the Lewis and Riesenfeld method [12] to obtain the corresponding Schrödinger states in terms of solutions of a second order ordinary differential equation. By constructing Gaussian wave packet states, they calculated the quantum dispersions, quantum correlations, and the uncertainty product for each mode of the quantized scalar field. Scalar fields have also been investigated in Ref. [13], where the authors construct the coherent states and established the existence of squeezed states.

The quantization of the electromagnetic field in the de Sitter spacetime could provide a better understanding of (time-dependent) cosmological models, e.g., the present distribution of the fluctuation spectrum of the electromagnetic fields can be traced back to the primordial spectrum of cosmological fluctuations.

In the context of the invariant method Pedrosa *et al.* [14] investigated the light propagation through time-dependent dielectric linear media in the absence of free charges and in a curved spacetime from a classical and a quantum point of view. They found that the light propagation present a remarkable similarity in both cases. From the classical study, they showed that the amplitude q of the potential vector ($\vec{A}$) is written in terms of the first- and second-order Bessel function for a propagation through a dielectric media with $\varepsilon(t) = \varepsilon_0 e^{\alpha t} (\alpha > 0)$. From the quantum perspective, they quantized the electromagnetic waves propagating in a material medium with $\varepsilon(t) = \varepsilon_0 e^{\alpha t}$, and q is written in terms of harmonic functions of exponentially decaying argument for a propagation in the de Sitter spacetime with the metric $a(t) = e^{H_0 t}$, where H_0 is the Hubble constant. They wrote the solutions of the Schrödinger equation in terms of ρ, which is solution of the Milne-Pinney equation [15, 16]. As a result they claim that for a problem with curved spacetime the electromagnetic field quantization can also be carried out following the same steps used for the time-varying dielectric case. Moreover, as in the classical case, the quantum behavior of the electromagnetic field is purely oscillatory in the conformal time in the de Sitter spacetime.

The study of de Sitter spacetime becomes important if one considers a ΛCDM model for the cosmos. In the present era, the universe can approximately be described by a de Sitter spacetime, where the matter decay with the volume and the cosmological constant is kept constant. For $t \gg H_0^{-1}$, the universe tends to be fully described by the de Sitter model. Here, we revisit the problem of quantizing the electromagnetic field in the de Sitter space as previously considered in Ref.[14]. We present an exact solution of the Schrödinger equation, construct the correspondent coherent states, and calculate various average values. The paper is organized as follows. In Sec. II we show how to obtain the correspondence between an electromagnetic field placed in a de Sitter background and a time-dependent harmonic oscillator. In Sec. III, we use the Lewis and Riesenfeld [12] method to obtain the solutions of the related Schrödinger equation, construct the coherent states, calculate the correlations between position and momentum and the uncertainty product. Section IV summarizes the results.

DECOMPOSITION OF THE VECTOR FIELD

Consider the electromagnetic field in a de Sitter space time. The action for the Gauge field is given by

$$S = -\frac{1}{4}\int d^4x \, \sqrt{-g} g^{\mu\nu} g^{\alpha\beta} F_{\mu\alpha} F_{\nu\beta}, \tag{1}$$

where the metric $ds^2 = -dt^2 + e^{2H_0 t} dx^2$ is diagonal and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, leading to

$$\begin{aligned} g^{\mu\nu} g^{\alpha\beta} F_{\mu\alpha} F_{\nu\beta} = & -2\delta^{ij} e^{-2H_0 t} F_{0i} F_{0j} + \\ & e^{-4H_0 t} \delta^{kl} \delta^{ij} F_{ki} F_{lj}. \end{aligned} \tag{2}$$

Using the Gauge conditions

$$A_0 = 0, \qquad \nabla \cdot \vec{A} = 0 \tag{3}$$

we obtain

$$S = -\frac{1}{2}\int d^4x e^{3H_0 t}(-e^{-2H_0 t}(\dot{A}_i)^2 + e^{-4H_0 t}(\partial_i A_j)^2) \tag{4}$$

where the dot denotes the time derivative. Now we decompose the field as

$$\vec{A}(\vec{x},t) = \sum_{\lambda=1,2} \int \frac{d^3k}{(2\pi)^3} \hat{\epsilon}^{(\lambda)} \left[u_k^{(\lambda)}(t) e^{i\vec{k}\cdot\vec{x}} + u_k^{(\lambda)*}(t) e^{-i\vec{k}\cdot\vec{x}} \right] \tag{5}$$

where $\hat{\epsilon}$ represents a unit vector satisfying $\vec{k} \cdot \hat{\epsilon}^{(\lambda)} = 0$ due to the Gauge condition and $\hat{\epsilon}^{(\lambda')} \cdot \hat{\epsilon}^{(\lambda)} = \delta^{\lambda'\lambda}$, and $\lambda = 1, 2$ stands for the two polarizations. By separating $u_k^{(\lambda)}$ into real and imaginary parts

$$u_k^{(\lambda)} = \frac{1}{\sqrt{2}} \sum_\lambda (q_1^\lambda + i q_2^\lambda) \tag{6}$$

the action reads

$$S = -\frac{1}{2}\int dt \int \frac{d^3k}{(2\pi)^3} \sum_{\lambda,i=1}^{2} -e^{H_0 t}\left[\dot{q}_{ki\lambda}^2 + k^2 q_{ki\lambda}^2\right], \tag{7}$$

where $i = 1, 2$ labels the real and imaginary parts of $u_k^{(\lambda)}$. From Eq.(7) we obtain the Hamiltonian for the electromagnetic field

$$H_{\lambda ik} = \frac{e^{-H_0 t}}{2}(p_{\lambda ik}^2 + k^2 q_{\lambda ik}^2), \tag{8}$$

where

$$p_{\lambda ik} = \frac{\partial L}{\partial q_{\lambda ik}} = e^{H_0 t} \dot{q}_{\lambda ik}, \tag{9}$$

with p being the conjugate momentum. The classical equation of motion for the qth mode reads

$$\ddot{q}_{\lambda ik} + H_0 \dot{q}_{\lambda ik} + \omega^2 q_{\lambda ik}^2 = 0. \tag{10}$$

Next, consider the classical harmonic oscillator with time-dependent mass and frequency given by the Hamiltonian

$$H(t) = \frac{p^2}{m(t)} + \frac{1}{2} m(t) \omega^2(t) q^2, \tag{11}$$

where $[q, p] = i\hbar$. The equation of motion reads

$$\ddot{q} + \frac{\dot{m}(t)}{m(t)} \dot{q} + \omega^2(t) q = 0, \tag{12}$$

which is similar to Eq. (10) if one considers that each mode of the electromagnetic field corresponds to the time-dependent harmonic oscillator with $m(t) = e^{H_0 t}$ and $\omega = k e^{-H_0 t}$.

QUANTIZATION OF ELECTROMAGNETIC FIELDS IN THE DE SITTER SPACETIME

Consider a time-dependent harmonic oscillator described by Eq. (11). It is well known that an invariant for Eq. (11) is given by [17]

$$I = \frac{1}{2}\left[\left(\frac{q}{\rho}\right)^2 + (\rho p - m\dot{\rho} q)^2\right] \tag{13}$$

where $q(t)$ sastisfy Eq.(12) and $\rho(t)$ satisfies the generalized Milne-Pinney equation [15**?** , 16]

$$\ddot{\rho} + \gamma(t)\dot{\rho} + \omega^2(t)\rho = \frac{1}{m^2(t)\rho^3}. \tag{14}$$

The invariant $I(t)$ satisfies the equation

$$\frac{dI}{dt} = \frac{\partial I}{\partial t} + \frac{1}{i\hbar}[I, H] = 0 \tag{15}$$

and can be considered hermitian if we choose only the real solutions of Eq. (14). Its eingenfunctions, $\phi_n(q,t)$, are assumed to form a complete orthonormal set with time-independent discret eigenvalues, λ_n. Thus

$$I\phi_n(q,t) = \lambda_n \phi_n(q,t) \tag{16}$$

with $\langle \phi_n, \phi_{n'} \rangle = \delta_{nn'}$. Taking he Schrödinger equation (SE)

$$i\hbar \frac{\partial \psi(q,t)}{\partial t} = H(t)\psi(q,t) \tag{17}$$

where $H(t)$ is given by Eq. (11) with $p = -i\hbar\frac{\partial}{\partial q}$. Lewis and Riesenfeld [12] showed that the solution $\psi_n(q,t)$ of the SE (see Eq.(17)) is related to the functions $\phi_n(q,t)$ by

$$\psi_n(q,t) = e^{i\theta_n(t)} \phi_n(q,t) \tag{18}$$

where the phase functions $\theta_n(t)$ satisfy the equation

$$\hbar\frac{d\theta_n(t)}{dt} = \left\langle \phi_n(q,t) \left| i\hbar\frac{\partial}{\partial t} - H(t) \right| \phi_n(q,t) \right\rangle. \tag{19}$$

The general solution of the SE may be written as

$$\psi_n(q,t) = \sum_n c_n e^{i\theta_n(t)}\phi_n(q,t) \tag{20}$$

where c_n are time-independent coefficients. Now, using an unitary transformation and following the steps as in Ref. [17] we find

$$\begin{aligned}\psi_n(q,t) =& e^{i\theta_n(t)}\left(\frac{1}{\pi^{1/2}\hbar^{1/2}n!2^n\rho}\right)^{1/2}\times\\ &\exp\left\{\frac{im(t)}{2\hbar}\left[\frac{\dot\rho}{2\hbar}+\frac{i}{m(t)\rho^2(t)}\right]q^2\right\}\times\\ &H_n\left(\frac{1}{\sqrt{\hbar}}\frac{q}{\rho}\right)\end{aligned} \tag{21}$$

where

$$\theta_n(t) = -(n+\frac{1}{2})\int_{t_0}^{t}\frac{1}{m(t')\rho^2}dt', \tag{22}$$

$\lambda_n = (n+\frac{1}{2})\hbar$, and $H_n$is the Hermite polynomial of order n.

For $m(t) = e^{H_0 t}$ and $\omega(t) = ke^{-H_0 t}$ the auxiliary Equation (14) reads

$$\ddot\rho + H_0\dot\rho + k^2 e^{-2H_0 t}\rho = \frac{e^{-2H_0 t}}{\rho^3}. \tag{23}$$

Following the procedure described in Ref. [15**?** , 16], we find $\rho = 1/k^{\frac{1}{2}}$, and from Eqs. (22) and (21) we have

$$\begin{aligned}\psi_n(q,t) =& \left(\frac{k}{\pi\hbar n!^2 2^n}\right)^{\frac{1}{4}} e^{-\frac{ik}{H_0}(n+\frac{1}{2})(e^{-H_0 t_0}-e^{-H_0 t})}\times\\ &\exp\left(-\frac{kq^2}{2\hbar}\right)H_n\left(\sqrt{\frac{k}{\hbar}}q\right)\end{aligned} \tag{24}$$

which, except for the phase factor, is similar to the well-known wave function for the time-independent harmonic oscillator.

The coherent states for the time-dependent harmonic oscillator (see Eq. (11)) are constructed as follows [18]. Consider the time-dependent creation $a^\dagger(t)$ and annihilation $a(t)$ operators defined as

$$\begin{aligned}a^\dagger(t) &= \frac{1}{\sqrt{2\hbar}}\left[\frac{q}{\rho} - i(\rho p - m(t)\dot\rho q)\right],\\ a(t) &= \frac{1}{\sqrt{2\hbar}}\left[\frac{q}{\rho} + i(\rho p - m(t)\dot\rho q)\right],\end{aligned} \tag{25}$$

where $[a^\dagger(t), a(t)] = 1$. In terms of $a(t)$ and $a^\dagger(t)$ the invariant I(see Eq. (13)) can be written as

$$I = \hbar(a(t)a^\dagger(t) + 1/2). \tag{26}$$

Let $|n,t\rangle$ be the eigenstates of I. Therefore the following relations hold

$$a(t)|n,t\rangle = \sqrt{n}|n-1,t\rangle, \tag{27}$$

$$a^\dagger(t)|n,t\rangle = \sqrt{(n+1)}|n+1,t\rangle, \tag{28}$$

$$I|n,t\rangle = \hbar(n+1/2)|n,t\rangle. \tag{29}$$

Since the coherent states for I can be easily constructed, the coherent states for the time-dependent harmonic oscillator are straightforwardly obtained:

$$|\alpha,t\rangle = e^{-\frac{|\alpha|^2}{2}}\sum_{n=0}^{\infty}\frac{\alpha}{n!^{1/2}}e^{i\theta_n(t)}|n,t\rangle, \tag{30}$$

where θ_n is given by (22), and the complex number $\alpha(t)$ satisfies the eigenvalue equation

$$a(t)|\alpha,t\rangle = \alpha(t)|\alpha,t\rangle, \tag{31}$$

with

$$\alpha(t) = \alpha(t_0)e^{2i\theta_0(t)} \tag{32}$$

and

$$\theta_0(t) = -\frac{1}{2}\int_{t_0}^{t}\frac{dt'}{m(t')\rho^2(t')}. \tag{33}$$

The fluctuations in q (Δq), p (Δp), and in the uncertainty product $\Delta q\Delta p$ of the coherent state $|\alpha,t\rangle$ are

$$\Delta q_\alpha = \sqrt{\frac{\hbar}{2}}\rho, \tag{34}$$

$$\Delta p_\alpha = \sqrt{\frac{\hbar}{2}}\frac{1}{\rho}(1+m^2(t)\dot\rho^2\rho^2)^{1/2}, \tag{35}$$

and

$$\Delta q_\alpha\Delta p_\alpha = \frac{\hbar}{2}(1+m^2(t)\dot\rho^2\rho^2)^{1/2}, \tag{36}$$

respectively.

For $\rho = 1/k^{1/2}$, we have $m(t)\rho\dot\rho = 0$ and as a consequence $\Delta q_\alpha\Delta p_\alpha$ is minimum $(\hbar/2)$. In this case, the coherent states $|\alpha,t\rangle$ are "true" minimum-uncertainty states. Otherwise, if $m(t)\rho\dot\rho \neq 0$, the states $|\alpha,t\rangle$ are not minimum-uncertainty states and are equivalent to the well-known squeezed states, as pointed out in Ref. [19]. This is an interesting result since the minimum uncertainty product is assumed to be satisfied only for time-independent harmonic oscillator, unless the solution of Eq.(14) is a constant.

Next, let us analyze the $\langle q\rangle_\alpha$ and $\langle p\rangle_\alpha$ time evolution and its phase diagram. By setting $\alpha(t_0) = u + iv$ we find

$$\Delta q_\alpha = \sqrt{2\hbar}\rho\,[u\cos(2\theta(t)) - \nu\sin(2\theta_0(t))] \tag{37}$$

and

$$\Delta p_\alpha = \sqrt{2\hbar}\left[\left(\frac{\nu}{\rho} + um\dot{\rho}\right)\cos(2\theta_0(t)) + \left(\frac{u}{\rho} - \nu m\dot{\rho}\right)\sin(2\theta_0(t))\right]. \tag{38}$$

The constants u and v are determined from the initial conditions $\langle q(t_0)\rangle_\alpha = q_0$ and $\langle p(t_0)\rangle_\alpha = p_0 = m(t_0)v_0$. Then we find

$$\langle q\rangle_\alpha = q_0 cos\left[\frac{k}{H_0}(e^{-H_0t} - e^{-H_0t_0})\right] - \frac{p_0}{k} sin\left[\frac{k}{H_0}(e^{-H_0t} - e^{-H_0t_0})\right], \tag{39}$$

and

$$\langle p\rangle_\alpha = p_0 cos\left[\frac{k}{H_0}(e^{-H_0t} - e^{-H_0t_0})\right] + kq_0 sin\left[\frac{k}{H_0}(e^{-H_0t} - e^{-H_0t_0})\right]. \tag{40}$$



FIG. 1. (a) The time evolution of $\langle q\rangle$, (b) Time evolution of $\langle p\rangle$, and (c) The phase space. We considered $k = 10^{-8}$ in all panels

Figures 1(a) and 1(b) show the time dependence of $\langle q\rangle_\alpha$ and $\langle p\rangle_\alpha$. Since we want to work in a de Sitter universe, we used the values: $t_0 \sim H_0^{-1} = 10^{18}$, $k = 10^{-8}$, $q_0 = 1.0$, and $p_0 = 1.0$. From Fig. 1(a) one can see that the system oscillates back and forth between the classical turning points with an increasing period and constant amplitude. For $t \approx 24H_o$, q and p reach their asymptotic values given in Eqs. (41) and (42).The phase diagram is shown in Fig. 1(c). Even though the dissapative nature of the system (the total energy ($E \approx e^{-H_0t}$)), it behaves like the usual time-independent harmonic oscillator (E =constant). This can be seen from the relation $B = \sqrt{2E/\lambda}$, where B is the amplitude of motion and λ the spring constant. Since $\lambda = k^2e^{-H_0t}$ and $E \approx e^{-H_0t}$, B is a constant. As t increases, the oscillation frequency $\omega(t)$ decreases while the oscillation period increases leading to the behavior shown in Fig. 1(a). It is also worth noticing that for $t >> (1/H_0)$, the oscillations disappear and the motion of both $\langle q\rangle_\alpha$ and $\langle p\rangle_\alpha$ tends to be a constant. This kind of behavior is expected in an eternally expanding universe.Therefore, in the above figure we consider our time in the range $t_0 = 10^{18} < t$ to illustrate the time evolution in this model. As said in the introduction, for the ΛCDM model, our universe gets closer and closer to de Sitter, this justifies the inclusion of large times ($t \to \infty$) in the figure. In Fig. 2 we show the behavior of the generalized coordinates and the phase space for a smaller value of the wave vector $k = 10^{-4}$. From this we can see that the modes with shorter wavelength oscillates for a longer time. This is expected since as the universe expand, the wavelengths are stretched and eventually disappear.

In the limit when $t \to \infty$, $\langle q\rangle_\alpha$ and $\langle p\rangle_\alpha$ are expressed as

$$\langle q\rangle_\alpha = cos\left(\frac{k}{H_0}e^{-H_0t_0}\right) + \frac{1}{k}sin\left(\frac{k}{H_0}e^{-H_0t_0}\right), \tag{41}$$

and

$$\langle p\rangle_\alpha = cos\left(\frac{k}{H_0}e^{-H_0t_0}\right) - ksin\left(\frac{k}{H_0}e^{-H_0t_0}\right). \tag{42}$$

From the expressions we see that the coordinates depends explicitly on k. We show in Fig 3 the behavior of the system when t goes to infinity. The $\langle q\rangle$ and $\langle p\rangle$ dependence on k is shown in panels a and b of the figure, while the phase space is shown in Fig. 3(c). In this Figure and the above expressions we see that for large values of k the $\langle p\rangle$ axis is more stretched while for small values this is so for the $\langle q\rangle$ axis.

Pedrosa *et al.* [19] have combined linear invariants and the LR method to obtain the exact wave function



FIG. 2. (a) Time evolution of $\langle q\rangle$(b) Time evolution of $\langle p\rangle$and (c) Phase space for $k = 10^{-4}$



FIG. 3. For $t \to \infty$: (a)k dependence of $\langle q\rangle$ (b)k dependence of $\langle p\rangle$ and (c) k dependence of phase space.

for a particle trapped by oscillating fields, written in terms of Mathieu functions. They calculated $\Delta q\Delta p$ and the quantum correlation between q and p, defined by $c_{1,1} = \frac{1}{2}\langle(qp + pq)\rangle - \langle q\rangle\langle p\rangle$. They are related through the equation

$$\Delta q\Delta p = \frac{\hbar}{2}\sqrt{1 + \left(\frac{2}{\hbar}c_{1,1}\right)^2}, \tag{43}$$

which shows that $\Delta q\Delta p$ is minimum whenever $c_{1,1} = 0$, as it happens for $c_{1,1}$ calculated in the coherent state $|\alpha, t\rangle$, *i.e.*, $(c_{1,1})_\alpha = 0$. The fact that $c_{1,1} = 0$ does not mean that q and p are uncorrelated. In order to verify the correlation between q and p one should study the function $c_{n,m} = \frac{1}{2}\langle(q^n p^m + p^m q^n)\rangle - \langle q^n\rangle\langle p^m\rangle$. For the coherent state $|\alpha, t\rangle$, we find that $(c_{2,2})_\alpha = -\hbar^2/2$, indicating that q and p even assumed as "classical" quantities are correlated.

The uncertainty product and correlations in the state ψ_n are more easily calculated using the relation $|\psi_n(q, t)\rangle = e^{i\theta_n(t)}|n, t\rangle$. They are given by

$$\Delta q_{\psi_n}\Delta p_{\psi_n} = \left(n + \frac{1}{2}\right)\hbar, \tag{44}$$

$$(c_{1,1})_{\psi_n} = 0, \tag{45}$$

and

$$(c_{2,2})_{\psi_n} = -\left(n^2 + n + \frac{1}{2}\right)\hbar^2. \tag{46}$$

We noticed that Eq. (37) is satisfied for $n = 0$. In this case ψ_0 is given by

$$\psi_0(q, t) = \left(\frac{k}{\pi\hbar}\right)^{\frac{1}{4}} e^{-\frac{ik}{2H_0}e^{-H_0 t}} \exp\left(-\frac{kq^2}{2\hbar}\right) H_0\left(\sqrt{\frac{k}{\hbar}}q\right) \tag{47}$$

in the coordinate representation of the coherent state.

CONCLUDING REMARKS

In this paper we used the Lewis and Riesenfeld method to obtain the time-dependent Schrödinger states emerging from the quantization of the electromagnetic field in the de Sitter spacetime. It is well-known that a challenge in obtaining the exact solution (see Eq. (21)) for the SE with H given in Eq. (11), is the solution of the Milne-Pinney (in terms of the universal scale factor of universe $a(t)$) equation

$$\ddot{\rho} + \frac{\dot{a}}{a}\dot{\rho} + \omega^2(t)\rho = \frac{1}{a^2(t)\rho^3}.$$

For a given $a(t)$, different eras of universe can be investigated. For the de Sitter universe $a(t) = e^{H_0 t}$, and $\rho{=}\frac{1}{k^{1/2}}$, and the solution for Ψ_n corresponds, except for the phase factor, to the well known wave functions of the time-independent harmonic oscillator.

We have constructed the "true" coherent states,$|\alpha, t\rangle$, whose coordinate representation is given by Eq. (47). We verified that Eq. (43) holds for $|\alpha, t\rangle$. We calculated the quantum fluctuations in the coordinate and momentum as well as the quantum correlations between the coordinate and momentum in the state $\Psi_n(q, t)$. We calculated the quantum fluctuations in the coordinate and momentum as well as the quantum correlations between the coordinate and momentum in the state $\Psi_n(q,t)$.

We analyzed the time behavior of q and p, as well as the phase diagram q and p (see Figs.1(a)-(c) and 2(a)-(c)). The phase diagram indicates that the properties of the light propagation through the de Sitter spacetime can be analyzed in terms of the well known time-independent harmonic oscillator in real time.

At last, we would like to point out that the procedure described can be used to trace the present properties of the quantum electromagnetic field back to the recombination era. This would be a much more interesting phenomenological result.We are currently investigating the properties of the quantum electromagnetic field for $a(t) \sim t^{\frac{1}{2}}$ and $a(t) \sim t^{\frac{2}{3}}$ to give a more complete description of the cosmological evolution of the quantized electromagnetic field.

ACKNOWLEDGMENTS

The authors would like to thank Jailson S. Alcaniz for useful discussions. We would also like to thank: The Goethe-Institut Berlin for the hospitality. We acknowledge the financial support provided by Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP), the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and FUNCAP/CNPq/PRONEX.

---

* geovamaciel@gmail.com
† ildeguedes@gmail.com
‡ renan@fisica.ufc.br
§ rai@fisica.ufc.br

[1] N. D. Birrell and P. C. W. Davies, *Cambridge, Uk: Univ. Pr. (1982) 340p*
[2] J. Polchinski, *Cambridge, UK: Univ. Pr. (1998) 402 p*
[3] J. Polchinski, *Cambridge, UK: Univ. Pr. (1998) 531 p*
[4] N. Berkovits, JHEP **0004**, 018 (2000) [arXiv:hep-th/0001035].
[5] S. W. Hawking, Nature **248**, 30 (1974).
[6] L. C. B. Crispino, A. Higuchi and G. E. A. Matsas, Rev. Mod. Phys. **80**, 787 (2008) [arXiv:0710.5373 [gr-qc]].
[7] A. A. Saharian and T. A. Vardanyan, Class. Quant. Grav. **26**, 195004 (2009) [arXiv:0907.1149 [hep-th]].
[8] L. Parker, Phys. Rev. Lett. **21**, 562 (1968).
[9] L. Parker, Phys. Rev. **183**, 1057 (1969).
[10] L. Parker, Phys. Rev. D **3**, 346 (1971) [Erratum-ibid. D **3**, 2546 (1971)].
[11] C. E. F. Lopes, I. A. Pedrosa, C. Furtado and A. M. de M. Carvalho, J. Math. Phys., 50, 083511 (2009).
[12] H. R. Lewis and W. B. Riesenfeld, J. Math. Phys. **10**, 1458 (1969).
[13] A.M.de M. Carvalho et al Phys. Rev D 70, 123523 (2004)
[14] I. A. Pedrosa, Claudio Furtado and Alexandre Rosas, Europhys. Lett. 94, 30002 (2011).
[15] E. W. Milne, Phys. Rev. 35 (1930) 863.
[16] E. Pinney, Proc. Am. Math. Soc. 1 (1950) 681.
[17] J. F. Cariñena, J. de Lucas, Int. J. Geom. Methods Mod. Phys., 6:4 (2009), 683–699.
[18] I. A. Pedrosa, Phys. Rev. A 55, 3219 (1997).
[19] I. A. Pedrosa, A. Rosas and I. Guedes, J. Phys Gen. 38, 7757 (2005).